

San Francisco | 15-19 December 2014

Water in the Cratonic Mantle: Insights from FTIR Data on Lac de Gras Xenoliths (Slave Craton, Canada)

Anne H Peslier¹, Alan D Brandon², Lillian Aurora Schaffer², Suzanne Yvette O'Reilly³, William L Griffin⁴, Richard V Morris⁵, Trevor G. Graff¹ and David G Agresti⁶, (1)Jacobs Technology, NASA Johnson Space Center, Houston, TX, United States, (2)University of Houston, Houston, TX, United States, (3)Macquarie University, Sydney, Australia, (4)Macquarie University, Department of Earth and Planetary Sciences, Sydney, Australia, (5)NASA Johnson Space Center, Houston, TX, United States, (6)University of Alabama at Birmingham, Birmingham, AL, United States

Abstract:

The mantle lithosphere beneath the cratonic part of continents is the deepest (> 200 km) and oldest (>2-3 Ga) on Earth, remaining a conundrum as to how these cratonic roots could have resisted delamination by asthenospheric convection over time. Water, or trace H incorporated in mineral defects, could be a key player in the evolution of continental lithosphere because it influences melting and rheology of the mantle [e.g., 1]. Mantle xenoliths from the Lac de Gras kimberlite in the Slave craton [2] were analyzed by FTIR. The cratonic mantle beneath Lac de Gras is stratified with shallow (<145 km) oxidized ultradepleted peridotites and pyroxenites with evidence for carbonatitic metasomatism, underlain by reduced and less depleted peridotites metasomatized by kimberlite melts [3,4]. Peridotites analyzed so far have H₂O contents in ppm weight of 7-100 in their olivines, 58 to 255 in their orthopyroxenes (opx), 11 to 84 in their garnet, and 139 in one clinopyroxene. A pyroxenite contains 58 ppm H₂O in opx and 5 ppm H₂O in its olivine and garnet. Olivine and garnet from the deep peridotites have a range of water contents extending to higher values than those from the shallow ones. The FTIR spectra of olivines from the shallow samples have more prominent Group II OH bands compared to the olivines from the deep samples, consistent with a more oxidized mantle environment [5]. The range of olivine water content is similar to that observed in Kaapvaal craton peridotites at the same depths (129-184 km [1]) but does not extend to as high values as those from Udachnaya (Siberian craton [6]). The Slave, Kaapvaal and Siberian cratons will be compared in terms of water content distribution, controls and role in cratonic root longevity.

- [1] Peslier et al. 2010 Nature 467, 78-81.
- [2] Aulbach et al. 2007 CMP 154, 409-427.
- [3] Creighton et al. 2009 CMP 157, 491-504.
- [4] Aulbach et al. 2013 CG 352, 153-169.
- [5] Bai & Kohlstedt 1993 PCM 19, 460-471.
- [6] Doucet et al. 2014 GCA 137, 159-187.

Back to: Volatiles and Earth's mantle processes

<< Previous Abstract | Next Abstract >>

SEARCH KEYWORDS
SEARCH SESSIONS
SEARCH CONVENER/AUTHOR NAME
LOGIN

AGU.org | About AGU | About the AGU Fall Meeting | AGU Meetings | FAQs | Sitemap |© 2014 American Geophysical Union. All Rights Reserved.

AGU galvanizes a community of Earth and space scientists that collaboratively advances and communicates science and its power to ensure a sustainable future.

2 of 3 8/20/14 1:12 PM

3 of 3 8/20/14 1:12 PM